



## Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland

Alexandre Buttler, B.J.M. Robroek, Fatima Laggoun-Défarge, Vincent E. J. Jassey, Cédric Pochelon, Gregory Bernard, Frédéric Delarue, Sébastien Gogo, Pierre Mariotte, Edward A. D. Mitchell, et al.

### ► To cite this version:

Alexandre Buttler, B.J.M. Robroek, Fatima Laggoun-Défarge, Vincent E. J. Jassey, Cédric Pochelon, et al.. Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland. *Journal of Vegetation Science*, 2015, 26 (5), pp.964-974. 10.1111/jvs.12296 . insu-01163362

**HAL Id: insu-01163362**

**<https://hal-insu.archives-ouvertes.fr/insu-01163362>**

Submitted on 12 Jun 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Journal of Vegetation Science

## Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland

Journal:	<i>Journal of Vegetation Science</i>
Manuscript ID:	JVS-RA-02979.R1
Manuscript Type:	Research article
Date Submitted by the Author:	n/a
Complete List of Authors:	Buttler, Alexandre; EPFL-WSL, Laboratory of Ecological Systems - ECOS Robroek, Bjorn; Utrecht University, Institute of Environmental Biology; EPFL-WSL, Laboratory of Ecological Systems - ECOS Laggoun, Fatima; CNRS/INSU, ISTO, UMR 7327 Jassey, Vincent; EPFL-WSL, Laboratory of Ecological Systems - ECOS Pochelon, Cédric; EPFL-WSL, Laboratory of Ecological Systems - ECOS Bernard, Gregory; Fédération des Conservatoires d'espaces naturels, Pôle-Relais Tourbières Delarue, Frédéric; Muséum National d'Histoire Naturelle, Laboratoire de Minéralogie et de Cosmochimie du Muséum, UMR 7202 CNRS Gogo, Sébastien; CNRS/INSU, ISTO, UMR 7327 Mariotte, Pierre; EPFL-WSL, Laboratory of Ecological Systems - ECOS Mitchel, Edward; Université de Neuchâtel, Laboratory of Soil Biology, Institute of Biology Bragazza, Luca; EPFL-WSL, Laboratory of Ecological Systems - ECOS
Keywords:	climate change, vegetation, Sphagnum, OTC, temperature, drought, NDVI, point-intercept survey

Research article

Estimated size: 6617 words (8 pages), figures 2 pages, total 10 pages

**Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland**

**Running head: Experimental warming and plant responses**

Alexandre Buttler, Bjorn J.M. Robroek, Fatima Laggoun-Défarge, Vincent E.J. Jassey, Cédric Pochelon, Gregory Bernard, Frédéric Delarue, Sébastien Gogo, Pierre Mariotte, Edward A.D. Mitchell & Luca Bragazza

Buttler, A. (Corresponding author, [alexandre.buttler@epfl.ch](mailto:alexandre.buttler@epfl.ch))<sup>1, 2, 3, 12</sup>

Robroek, B.J.M. ([bjorn.robroek@epfl.ch](mailto:bjorn.robroek@epfl.ch))<sup>1, 2, 11</sup>

Laggoun-Defarge, F. ([Fatima.Laggoun-Defarge@univ-orleans.fr](mailto:Fatima.Laggoun-Defarge@univ-orleans.fr))<sup>4, 5, 6</sup>

Jassey, V.E.J. ([vincent.jassey@epfl.ch](mailto:vincent.jassey@epfl.ch))<sup>1, 2, 3</sup>

Pochelon, C. ([cedric.pochelon@bluewin.ch](mailto:cedric.pochelon@bluewin.ch))<sup>1, 2</sup>

Bernard, G. ([gregory.bernard@reseau-cen.org](mailto:gregory.bernard@reseau-cen.org))<sup>1, 2, 14</sup>

Delarue, F. ([fdelarue@mnhn.fr](mailto:fdelarue@mnhn.fr))<sup>4, 5, 6, 7</sup>

Gogo, S. ([Sebastien.Gogo@univ-orleans.fr](mailto:Sebastien.Gogo@univ-orleans.fr))<sup>4, 5, 6</sup>

Mariotte, P. ([pierre.mariotte@hotmail.com](mailto:pierre.mariotte@hotmail.com))<sup>1, 2, 12, 13</sup>

Mitchell, E.A.D. ([edward.mitchell@unine.ch](mailto:edward.mitchell@unine.ch))<sup>9, 10</sup>

Bragazza, L. ([luca.bragazza@wsl.ch](mailto:luca.bragazza@wsl.ch))<sup>1, 2, 8</sup>

Author contributions: A.B. wrote the manuscript, A.B., G.B and C.P. collected the vegetation data, A.B., F.L.D, S.G. and E.M. conceived the experiment, B.R., V.J. and F.D. helped for vegetation and micrometeorological data analysis and all contributed to the interpretation of results and commented on the manuscript.

#### Addresses:

<sup>1</sup>Ecole Polytechnique Fédérale de Lausanne EPFL, School of Architecture, Civil and Environmental Engineering ENAC, Laboratory of ecological systems ECOS, Station 2, 1015 Lausanne, Switzerland.

<sup>2</sup>Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Site Lausanne, Station 2, 1015 Lausanne, Switzerland.

<sup>3</sup>Laboratoire de Chrono-Environnement, UMR CNRS 6249, UFR des Sciences et Techniques, 16 route de Gray, Université de Franche-Comté, F-25030 Besançon, France.

<sup>4</sup>Université d'Orléans, ISTO, UMR 7327, 45071 Orléans, France.

<sup>5</sup>CNRS/INSU, ISTO, UMR 7327, 45071 Orléans, France.

<sup>6</sup>BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France.

<sup>7</sup>Laboratoire de Minéralogie et de Cosmochimie du Muséum, UMR 7202 CNRS, Muséum National d'Histoire Naturelle, 57 rue Cuvier, 75231 Paris Cedex 05, France.

<sup>8</sup>University of Ferrara, Department of Life Science and Biotechnologies, Corso Ercole I d'Este 32, I-44121 Ferrara, Italy.

<sup>9</sup>Laboratory of Soil Biology, Institute of Biology, University of Neuchâtel, CH-2000 Neuchâtel, Switzerland.

<sup>10</sup>Jardin Botanique de Neuchâtel, Chemin du Perthuis-du-Sault 58, 2000 Neuchâtel, Switzerland.

<sup>11</sup>Ecology and Biodiversity, Institute of Environmental Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands.

<sup>12</sup>Department of Environmental Science, Policy and Management, Suding Lab, University of California Berkeley, 147 Hilgard Hall, Berkeley, CA 94720.

<sup>13</sup>University of Sydney, Centre for Carbon, Water and Food Biogeochemistry Group - 380 Werombi Road, Camden NSW 2570.

<sup>14</sup>Pôle-relais Tourbières, Fédération des Conservatoires d'espaces naturels, 7, rue Voirin, 25000 Besançon, France.

Keywords: climate change; vegetation, *Sphagnum*; OTC; temperature; drought; NDVI; point-intercept survey.

**ABSTRACT**

Question: A better understanding of the response of *Sphagnum* mosses and associated vascular plants to climate warming is relevant for predicting the carbon balance of peatlands in a warmer world. Open-top chambers (OTCs) have been used to investigate the effect on soil biogeochemical processes in peatlands, but little information is available on the effects of OTCs on microclimate conditions and the associated response of plant community. We aimed to understand how simulated warming and differences in soil moisture affect plant species cover.

Methods and location: We used OTCs to measure the effect of a near-ground temperature increase (+ 1.5 °C on average) on vegetation dynamics over five growing seasons (2008-2012) in a *Sphagnum*-dominated peatland (French Jura), in two adjacent microhabitats with different hydrological conditions – wet and dry. Microclimatic conditions and plant species abundance were monitored at peak biomass in years 1, 2, 3 and 5 and monthly during the plant-growing season in year 5.

Results: The response of plants to warming differed between vascular plants and bryophytes as well as among species within these groups and also varied in relation to soil moisture. *Andromeda polifolia* abundance responded positively to warming, while *Vaccinium oxycoccus* responded negatively and *Eriophorum vaginatum* showed a high resistance.

Conclusion: Depth of rooting of vascular plants appeared to control the response in plant abundance, while moss abundance depended on various other interacting factors, such as shading by the vascular plant community, precipitation and soil moisture.

## INTRODUCTION

Peatlands are long-term sinks of atmospheric carbon (C) due to the accumulation of peat (Yu 2011). These ecosystems develop in areas that are hydrologically and climatically characterized by a net surplus of water. As a consequence, prevailing soil anoxic conditions hamper microbial decomposition of organic matter (Holden 2005).

In northern hemisphere ombrotrophic (= rain fed) peatlands, bryophytes of the genus *Sphagnum* (peat mosses) are generally the dominant plant species. Peat mosses produce litter with antimicrobial properties, which hampers microbial decomposition (Hajek et al. 2011), leading to the build-up of peat. Further, *Sphagnum* mosses can affect the abundance and the performance of associated vascular plants (Dorrepaal et al. 2006) by regulating soil moisture, energy balance, soil temperature, and nutrient availability (Turetsky et al. 2012). Ultimately, the inter-specific competition between *Sphagnum* mosses and vascular plants controls peat accumulation because an increase of vascular plant abundance and in particular their shading effect can reduce *Sphagnum* growth (Heijmans et al. 2002; Bragazza et al. 2013). Changes in the competitive balance between plant species of different growth forms has been shown to alter the C balance in peatlands (Kuiper et al. 2014), potentially by changes in the litter quality, which ultimately affects microbial decomposition (Bragazza et al. 2007; Dorrepaal et al. 2007; Gogo et al. 2011).

Climate warming and associated extreme climatic events can be detrimental to *Sphagnum* mosses as, like all terrestrial bryophytes, these plants cannot actively control their water balance and

therefore rely on atmospheric precipitation to maintain suitable hydrological conditions (Robroek et al. 2007, Bragazza 2008; Nijp et al. 2014). Increased air temperature can increase evapotranspiration with direct negative effects on *Sphagnum* productivity (Gerdol 1995; Weltzin et al. 2001), but with positive effects on vascular plant growth by drying out the top soil and enhancing soil oxygenation (Weltzin et al. 2000; Bragazza et al. 2013), to which plants of different functional types might respond differently (Kuiper et al. 2014). A better understanding of the response of *Sphagnum* mosses and associated vascular plants to climate warming is then relevant for predicting the C balance of peatlands in a warmer world. Indeed, a warmer climate has been reported to alter plant species abundance in peatlands, in particular favoring ericaceous dwarf shrubs (Weltzin et al. 2003, Breeuwer et al. 2009), which will affect soil biogeochemical processes (Bragazza et al. 2013).

Under field conditions, simulation of climate warming can be obtained by using passive warming open-top chambers (OTCs) (Marion 1997). Although OTCs are recognized to increase temperature consistent with general circulation models, they also affect a set of microclimatological variables such as moisture, light, snow accumulation and wind speed (Bokhorst et al. 2013). In peatland studies, OTCs have been reliably used to investigate the effect on soil biogeochemical processes (Dorrepaal et al. 2009), but little information is available on the effects of OTCs on microclimate conditions and the associated response of plant species (Weltzin et al. 2003, Keuper et al. 2011; Delarue et al. 2014; Jassey et al. 2013).

In the present study we used OTCs to simulate increased soil and air temperature, and we monitored changes in vegetation cover in a *Sphagnum*-dominated peatland over five years of experimental warming. We aimed to understand how species abundance and plant community composition responded to simulated warming and associated changes in soil moisture. We hypothesized that plant responses will vary among functional types; and more specifically that 1) vascular plants will respond to warming according to their rooting depth, and 2) *Sphagnum* mosses

will have a more complex response depending on the interaction between micrometeorological conditions and abundance of vascular plants.

## MATERIAL AND METHODS

### *Study site*

The study was performed on an undisturbed, ombrotrophic *Sphagnum*-dominated peatland situated in the Jura Mountains (The Forbonnet peatland, France, 46°49'35''N, 6°10'20''E, 840 m a.s.l.). Climate is characterized by relatively cold winters (average temperature -1.4°C) and mild summers (average temperature 14.6°C), with annual mean temperature of 7.5°C and annual mean precipitation between 1300 and 1500 mm (Laggoun-Défarge et al. 2008).

Two areas (microhabitats) were selected based on their hydrological conditions. The first microhabitat (hereafter called 'wet') was a *Sphagnum*-dominated relatively flat area bordering a transitional poor fen, characterized by a moss cover dominated by *Sphagnum fallax*. Vascular plants such as *Eriophorum vaginatum* and *Andromeda polifolia* were recorded with low abundance, together with *Scheuchzeria palustris*, *Drosera rotundifolia* and *Vaccinium oxycoccus*. The second microhabitat (hereafter called 'dry') was adjacent to the wet microhabitat, but in a area with generally deeper water tables. The dry microhabitat was characterized by a pattern of hummocks dominated by *Sphagnum magellanicum*, *V. oxycoccus*, *E. vaginatum* and *Calluna vulgaris*, and lawns with *S. fallax*, *Carex rostrata* and *A. polifolia*. Tree (*Pinus rotondata*) encroachment occurs in this dry microhabitat.

### *Experimental design*

In April 2008, in both the dry and the wet microhabitats, six representative plots (3 × 3 m) were selected, which were then randomly allocated to either the control or the warming treatment



(Laggoun-Défarge et al., 2008). Warming treatments were obtained by using open-top chambers (OTCs), which consisted of transparent polycarbonate hexagonal chambers (50 cm high, 1.7 m top-width, 2.4 m base-width). OTCs allow quasi-natural transmittance of visible wavelengths, and minimize the transmittance of re-radiated infrared wavelengths (Marion et al., 1997). All OTCs were raised 10 cm above the peat surface to allow air circulation. The minimal (adjacent plots) and maximal (distant plots) distance between plots were *c.* 5 and 30 m, respectively.

#### *Environmental monitoring*

From November 2008 to November 2012, peat temperature (7 cm below the moss carpet) and air temperature (at 10 cm above the moss carpet) were monitored at 30-minute intervals using thermocouple probes connected to a data logger (CR-1000 Campbell). Sensors were repositioned as necessary to keep height and depth of measurement constant despite moss carpet accumulation. In 2012, these measurements were augmented with combined soil temperature/moisture sensors. In each microhabitat (wet/dry) and treatment (OTC/control) a randomly chosen plot was equipped with soil temperature-moisture probes (Decagon 5TM) at 3 cm and 5 cm depth. Furthermore, precipitation at the study site and water table depth in randomly chosen plot were continuously monitored by a meteorological station and sensors connected to the CR-1000 data logger.

#### *Plant species abundance*

At peak biomass (July or August, depending on the year) in 2008, 2009, 2010 and 2012, plant species abundance was measured using the point-intercept frequency method (Buttler 1992) in 50 × 50 cm subplots located in flat lawns. We used a Plexiglas frame with adjustable legs placed above a permanently marked subplot. A 20 holes ruler was moved along 20 different positions so as to obtain 400 measuring points on a regular grid. A metal pin (1 mm diameter) was lowered through each hole in the ruler and each contact of the pin with living vegetation and litter was recorded by

species until the pin reached the *Sphagnum* surface. This field technique could account for three-dimensional biomass distribution because, at each point, each plant that was intercepted was recorded. The relative frequency of each species was then calculated.

To allow analyses on seasonal dynamics of plant species cover, during the growing season 2012 (April - September), monthly pictures were taken from aforementioned subplots. In order to obtain high-resolution images, the subplots were divided into four  $25 \times 25$  cm quadrats. For the digital image analysis, a grid of 100 points was laid on each  $25 \times 25$  cm picture, after which the four grids were merged into a single 400 points grid-picture. Species overlaying the grid intersects were identified at 200% digital magnification. The relative frequency of each species was then calculated for each sampling date, as for the point-intercept method. Because we could not reliably distinguish *S. fallax* and *S. rubellum* in our point-intercept field measures and in the frequency analysis on photographs, we pooled these two species.

#### *Sphagnum height growth*

The cranked wire method (Clymo 1970) was used to measure *Sphagnum* growth. At the start of the 2011 and 2012 growing seasons (early April, after snow melt) cranked wires (5-10) were placed in each subplot. At the beginning of November, before snowfall, the length of the cranked wire above the moss surface was measured with a graduated pipe sliding down along the cranked wire and the average height of the *Sphagnum* capitula around the wire was calculated.

#### *Seasonality of plant community*

The normalized difference vegetation index (NDVI) was used as a phenological proxy and measured every fortnight between March and November 2012 using a portable spectroradiometer sensor. This measure is also used as an indirect measure of bulk biomass (Soudani et al. 2012). The sensor was maintained at 50 cm above the moss carpet, which allowed obtaining a NDVI signal of

the whole plant community, including the highest plants, while covering only the subplot surface. In each 50 × 50 cm subplot and for each campaign of sampling, 10 consecutive measures were taken according to a grid sampling design and then averaged to have a single NDVI value per plot. The NDVI measurements were performed under direct sunlight and with an umbrella to shadow the subplot.

# *Statistical analysis*

Differences in vegetation between years (i.e. 2008, 2009, 2010, 2012) were analyzed using relative species frequency data obtained from the point-intercept method in the field, whereas seasonal trend of plant species cover during the 2012 growing season was analyzed with species relative frequency data obtained from the photographic image analysis. Analysis at species level was done with Linear Mixed Models, with treatment (OTC vs. control) and years (or months for the seasonal data set in 2012) as fixed factors, and years (or months) nested in plots and blocks (wet and dry microhabitats) as random factors. The same model was also used to test the block effect. *Sphagnum* growth was analyzed with Linear Mixed Models, with treatment and block as fixed factors, and location of growth measures within plots given as random nested factors. The model was also tested for each block separately. NDVI was also analyzed with Linear Mixed Model, with treatment and block as fixed factors, and dates nested in plots as random factors. Random effects on the intercept were used to correct the inflation of the residual degrees of freedom that would occur if repeated measurements within sites would have been used as true replicates (Pinheiro & Bates 2000). In addition, for NDVI, separate ANOVA's were performed for each block and date. Pearson's correlation coefficients were calculated and tested between NDVI signals and species frequencies. Differences in monthly means of differences in daily mean air and soil temperature were assessed by ANOVA's. Comparisons of mean water level depth between blocks were done

with Student t-tests. Data were log transformed prior to analysis. All statistical analyses were performed in R (R Core Team, 2014).

## Results

### *Micrometeorology*

In general, over the period 2009-2012, monthly air temperature in the OTCs was 0.2-1.5 °C higher than in the control plots ( $p < 0.05$ ). Interestingly, OTCs rarely affected soil temperature, and when they did ( $p < 0.05$ ), the pattern was less clear, varying between a warming effect up to 0.3 °C in some cases, and a cooling effect up to -1 °C in some others (see also Delarue et al. 2011, 2014, 2015, Jassey et al. 2011). These effects of OTC on soil temperature were not related to the season, neither to the microhabitat. Soil moisture and temperature measured in 2012 with Decagon sensors in a subset of 4 plots (dry control, dry OTC, wet control and wet OTC) indicate that soil temperature was slightly higher in OTCs this particular year (mean daily differences between OTC and control plots for soil temperature was +0.6 °C in both microhabitat), and that moisture was slightly lower in OTC plots (mean daily differences between OTC and control plots for soil moisture were -0.011 and -0.067 (VWC) in dry and wet microhabitat, respectively).

Monthly cumulative rainfall during the period of *Sphagnum* growth measurements was higher in 2012 (2108 mm) than in 2011 (1376 mm). Depth of water level fluctuated over time, but highest water levels were recorded during spring 2012. Mean water level during the measurement period of *Sphagnum* growth was significantly higher in 2012 than in 2011 (Student t-test;  $p = 0.002$  in dry,  $p = 0.004$  in wet). Mean water level was also higher in the wet than the dry microhabitat ( $p = 0.004$ ).

### *Annual change in species abundance*

Considering all vascular plants, abundance (relative frequency) increased significantly with the OTC treatment (Figure 1). At species level, it appeared that with warming *A. polifolia* increased its

abundance consistently in wet and dry microhabitats, whereas the abundance of *V. oxycoccus* decreased. *D. rotundifolia* abundance responded marginally to the OTC treatment whereas *E. vaginatum* did not respond. Some other species responded contrastingly to the OTC treatment in the two microhabitats, for example *P. strictum* and *S. fallax+rubellum* increased their abundance in the dry microhabitat, whereas their abundance decreased or did not change in the wet microhabitat (treatment x block  $p=0.0367$  and  $0.0767$ , respectively). Litter increased in the wet microhabitat, and decreased in the dry microhabitat (treatment x block  $p=0.0001$ ). Significant changes occurred also over time, with a decrease of abundance for *V. oxycoccus* and *S. fallax+rubellum*, and an increase for *E. vaginatum* and litter.

There was also a strong microhabitat effect related to the dry and wet situations ( $p<0.05$ ), except for *Drosera rotundifolia* for which the microhabitat effect was marginally significant, and for *Carex rostrata*, *C. limosa*, *Calluna vulgaris* and *Pinus rotundata* for which there was no significant microhabitat effect. *Vaccinium oxycoccus*, *D. rotundifolia*, *Scheuchzeria palustris*, and *Sphagnum fallax+rubellum* were more abundant in the wet microhabitat, whereas species such as *Andromeda polifolia*, *Eriophorum vaginatum* and *Polytrichum strictum* were more frequent in the dry microhabitat, or exclusively found in the latter one (*S. magellanicum*).

#### Monthly change in species abundance

During the 2012 season, total vascular plant abundance and, more specifically *Andromeda polifolia* frequency increased significantly in the OTCs (Figure 2). *Vaccinium oxycoccus* abundance decreased in the wet microhabitat (treatment x block  $p=0.0034$ ) whereas that of *Sphagnum fallax+rubellum* declined in the wet microhabitat, but increased in the dry microhabitat (treatment x block  $p=0.008$ ). Litter increased in the wet microhabitat and decreased in the dry microhabitat (treatment x block  $p=0.0001$ ). Significant changes occurred also over the monthly course of 2012,

with a decrease in abundance for *S. magellanicum* (present only in the dry microhabitat) and litter, and an increase for *V. oxycoccus* and total vascular plants.

#### *Sphagnum* growth

The length increment of *Sphagnum* species differed between years, being higher in 2012 as compared to 2011 (Figure 3). In 2011, *Sphagnum* growth was significantly lower in the wet microhabitat, but warming did not affect the length increment. In 2012, there were significant microhabitat and treatment effects as well as a cross effect, with increased moss growth in the warmed plots of the dry microhabitat ( $p=0.017$ ), whereas no-significant changes were observed in the wet microhabitat.

#### Normalized Difference Vegetation Index

NDVI values taken under direct sunlight and in the shade were strongly correlated ( $R^2=0.83$ ), yet measurements at the start of the season (early May) were less correlated ( $R^2=0.49$ ) than measurements close to the peak of biomass (mid-June;  $R^2=0.82$ ). The precision of the measurements was calculated as the mean difference between each pairwise measurement. It gives an error of 1.6% under sun and 2.0% in the shade. As such, we only used NDVI data from sun measurements in further analyses.

Throughout the season, OTCs increased the vegetation's NDVI signal ( $p=0.001$ ), yet the effect of OTCs differed between wet and dry microhabitats (block x treatment  $p=0.002$ ). In the dry microhabitat, the NDVI signal in the OTC exceeded the signal in control plots throughout the season (Figure 4a). Such difference was mainly significant at the beginning of the growing season, and marginally different in summer. In the wet microhabitat, however, the difference between OTC and control plots was less obvious, and occurred during a more limited time in summer (Figure 4b).

NDVI was positively correlated to vascular plant abundance ( $r=0.85$ ,  $p<0.001$ ), and negatively with litter ( $r=-0.60$ ,  $p<0.001$ ). The correlation between NDVI and *Sphagnum* cover was negative ( $r=-0.24$ ,  $p<0.045$ ).

## Discussion

### *Effect of OTC on microclimate*

The observed OTC warming effect on near-ground air temperature was in the range of those found in other studies (e.g., Marion et al., 1997; Hollister & Webber, 2000; Dorrepaal et al. 2003; Sullivan et al., 2005, Walker et al. 2015). Yet, in accordance with Dabros et al. (2010), we observed that OTCs can reduce soil temperature, as a result of increased evapotranspiration (Delarue et al. 2011). Daily temperature fluctuations, minimum and maximum values and day or night means have all been shown to be affected by OTC warming (e.g. Marion et al. 1997; Dabros et al. 2010). A previous study at the same site revealed that daily soil temperature amplitudes were smaller but often significantly different between OTC's and control plots in winter months (Delarue et al. 2011). In summer months, amplitudes were larger, and even more so in OTCs as compared to control plots in April, May and June in the dry microhabitat (but differences were marginally or not significant), whilst an opposite trend appeared in the wet microhabitat. The additional Decagon sensors installed in 2012, which integrate more soil volume, might be more adapted to the very porous peat soil than tiny thermocouples. Despite the absence of replication, it is reasonable to recognize that our OTCs had a slight warming effect on soils for the largest part of time, yet this effect coincided with a slight reduction in soil volumetric water content as previously reported (Jassey et al. 2011)

### *Effect of OTCs on vegetation*

We showed that experimental warming differently influenced plant abundance (or frequency),

331 depending on the underlying hydrological conditions and on individual species identity. *A. polifolia*  
332 increased in abundance while *V. oxycoccus* decreased, and both *S. fallax+rubellum* and *P. strictum*  
333 responded positively to the OTC treatment in the dry microhabitat, and decreased or remained  
334 constant in the wet microhabitat. Among these species, the response of *A. polifolia* is interesting.  
335 This species has been shown to be sensitive to changes in environmental conditions such as  
336 snow cover and spring/summer warming (Aerts et al. 2006; Jassey et al. 2013; Robroek et al.  
337 2013). It allocates about 98% of the total biomass belowground, concentrated in the 0-15 cm  
338 peat layer, with the bulk at 2.5-7.5 cm, and a continuous rooting system from recent vegetation  
339 down into the peat (Wallén 1986). Roots of *A. polifolia* may reach a depth of 45 cm (Jacquemart  
340 1998) and fine roots form up to 24% of the total below ground biomass (Wallen 1986). *A. polifolia*  
341 might therefore be expected to have a somewhat deeper root system as compared to several other  
342 species enabling the species to sustain growth when soil dries out at the surface. Differently, *V.*  
343 *oxycoccus* has a shallow root system hardly penetrating more than a few centimeters below the  
344 living parts of the moss layer (Jacquemart 1998). With its shallow roots, this species relies on the  
345 water-conducting capacity of the *Sphagnum* mosses for its water supply (Malmer et al. 1994).  
346 Indeed, a decrease of *Sphagnum* water content in the OTC treatment has been demonstrated by  
347 Jassey et al. (2011). A subsequent lowering of water table will also reduce the growth rate of *V.*  
348 *oxycoccus* through water stress (Malmer et al. 1994, Jacquemart 1998). Coriaceous leaves of *A.*  
349 *polifolia*, their silvery-glaucous lower surface and their revolute margins, could indicate drought  
350 resistance. But anatomical features of *V. oxycoccus* such as thick epicuticular waxes and stomata on  
351 the abaxial surface only are considered similar to those of desert plant characteristic to a high ability  
352 to avoid drought (Jacquemart 1997). Drier soil might also promote ericoid mycorrhizal infection  
353 (Cullings 1996) and thus better access to nutrients in the soil, but this might equally apply to other  
354 Ericaceous present in the bog, such as *V. oxycoccus*.



355 An alternative explanation for the decrease of *V. oxycoccus* with OTC warming may come from  
356 direct nutrient resource competition (Malmer et al. 1994), in particular with *A. polifolia*. Because,  
357 however, OTC warming coincided with a decrease in moisture content, a direct effect of moisture  
358 limitation on the shallow rooting *V. oxycoccus* cannot be ruled out. The evidence of moisture  
359 limitation is supported by the lack of response to the OTC by *E. vaginatum*. This species is known  
360 to root deeply into the peat, up to 1.5 meters, with roots having highest elongation rates at low soil  
361 temperatures (Shaver & Billings 1977). Contrary to *A. polifolia* and *V. oxycoccus*, no mycorrhizal  
362 association has been reported for *E. vaginatum*. In warming experiment, no significant differences  
363 were reported for *E. vaginatum* root production between OTC's and control plots, albeit a tendency  
364 to higher production was observed (Sullivan & Welker 2005). We believe that the resistance of *E.*  
365 *vaginatum* is higher as compared to more shallow rooted species, and that an effect of warming has  
366 to be expected on a longer term, as soil warms up at lower depth. Weltzin et al. (2003) found that  
367 within bog and fen communities, different life forms and species can respond differently to  
368 warming and water table manipulation. They showed that warming may lead to an increase in  
369 abundance of woody plants, as does lowering water table as a result of changes in rates of  
370 evapotranspiration. In the peatlands studied by Weltzin et al. (2003), *Andromeda glaucophylla*  
371 responded mostly to the heat treatment, while other dominant shrubs (such as *Vaccinium oxycoccus*)  
372 did not, whereas graminoids decreased their cover. As observed by Weltzin et al. (2000), change in  
373 belowground biomass can be even more important. Our wet and dry sites showed similar trends for  
374 vascular plants, but not for mosses. The better growth of some mosses in warmed plots might be  
375 transient since, on the long run, mosses will be out-competed by the concomitant increase of  
376 *Andromeda* shrubs. Overall, these observations could point to a different evolution of such  
377 contrasted dry and wet situations, as for "bog" and "fen" like facies (Weltzin et al. 2003).

378 The increase of litter in the wet microhabitat and its decrease in the dry microhabitat under the  
379 OTC treatment does not follow the pattern of the abundance of total vascular plants, which could

indicate that litter abundance is mainly driven by the differential decomposition rate in the dry and wet microhabitats. Nevertheless, this relationship between warming and decomposition is not straightforward (Aerts 2006) and would deserve a proper litter decomposition experiment.

Moss length increment was also higher in the dry microhabitat in 2011, and was promoted by OTC warming in 2012 (Figure 3). In the dryer 2011 year, the higher cover of vascular plants, i.e. *Eriophorum* leaves in the dry microhabitat is the determinant factor for moss growth and might have had a facilitating effect on *S. fallax+rubellum* in limiting the drying out of the moss carpet (Grosvernier et al. 1994, Buttler et al. 1998, Walker et al. 2015). Instead, in the wetter year 2012, when water limitation was not critical and engendered an overall higher length increment, the warming effect by OTC's could promote further the moss growth in the dry microhabitat. Adaptive seasonal acclimation has been shown for *S. fallax*, as reflected by the maintenance of relatively high net photosynthetic rates to lower water contents and no response differences when plants were collected from different heights above the water table (Titus et al. 1983). In our study, we pooled *S. fallax* and *S. rubellum*, because we could not reliably distinguish these two species in our point-intercept field measures and in the frequency analysis on photographs. Nevertheless, some field sampling in the plots showed that *S. fallax* is by far the most frequent species and that *S. rubellum* occurs only on some very limited patches. Similarly to *S. fallax*, *S. rubellum* has been shown to adapt and maintain similar height increment at different experimental water table depths (Robroek et al. 2007). Therefore, shading may be critical for the development of these mosses, mediating soil moisture and near ground temperature, which both are direct drivers of growth. Our results show that shading can facilitate growth in dry years, but when wetter years are combined with warmer conditions, in our case the OTC effect, this shading effect on growth is out-ruled by temperature. Nevertheless, a better length increment of *Sphagnum* might also, on the long run, change the structure of the moss carpet and consequently its moisture holding and transport capacity (Dorrepaal et al. 2003). Therefore, when analyzing long term-effect of global change, not only the

change in species composition should be considered (Heijmans et al. 2008), but also the change of vegetation structure might be critical. In their work, Walker et al. (2015) mentioned a mismatch in the growth responses of different plant functional types to simultaneous climate and vegetation change. This was due to the complexity of interacting factors between dwarf-shrubs, bryophytes and graminoids, suggesting that some functional types reflecting a direct response to increased air temperature, some others an indirect one.

The evolution of aboveground biomass over the seasons was well reflected in the NDVI, which was also higher in OTC plots. Interestingly, NDVI was also better correlated to vascular plants than to mosses, which explains the more obvious difference in the dry plot, where vascular plants are more abundant. Indeed, when mosses dry out temporarily in dry periods, which can be observed in the wet microhabitat where the moss carpet is less protected from wind, reflectance is lowered. This differentiation between mosses and higher plants is critical in studying the carbon cycle using airborne remote sensing (Yan et al. 2014).

## Conclusion

Our study on the response of plant cover to experimental warming in two contrasting hydrological peatland microhabitats – wet and dry – shows that experimental warming by OTC interacts with soil wetness, resulting in divergent responses of vascular plants and bryophytes. For vascular plants, the depth of rooting seems the key in determining the response of plant cover, whereas for mosses, the growth is under the dependence of various other interacting factors, such as shading by shrubs and large statured vascular plants, which contributes further to regulate the microclimate conditions in the moss carpet.

## Acknowledgments

This work was funded as part of the PEATWARM initiative through an ANR (French National Agency for Research) grant (ANR-07-VUL-010). We acknowledge support from Labex VOLTAIRE (ANR-10-LABX-100-01) and grant PSPB-013/2010 from Switzerland through the Swiss Contribution to the enlarged European Union. P.M. acknowledges the support of the Swiss National Science Foundation (n<sup>o</sup> PBELP3 146538). The authors are also indebted to the Regional Scientific Council of Natural Heritage of the Franche-Comté Region for permitting access to Le Forbonnet site. We also thank M.-L. Toussaint for providing the meteorological data, E. Farrer and L. M. Hallet for statistical advices, and B. Corboz for technical assistance.

## References

- Aerts, R. 2006. The freezer defrosting: global warming and litter decomposition rates in cold biomes. *Journal of Ecology* 94: 713-724.
- Aerts, R., Cornelissen, H. & Dorrepaal, E. 2006. Plant performance in a warmer world: general responses of plants from cold, northern biomes and the importance of winter and spring events. *Plant Ecology* 182: 65-77.
- Bragazza, L., Siffi, C., Iacumin, P. & Gerdol, R. 2007. Mass loss and nutrient release during litter decay in peatland: the role of microbial adaptability to litter chemistry. *Soil Biology & Biochemistry* 39: 257-267.
- Bragazza, L. 2008. A climatic threshold triggers the die-off of peat mosses during an extreme heatwave. *Global Change Biology* 14: 2688-2695.
- Bragazza, L., Parisod, J., Buttler, A. & Bardgett, R.D. 2013. Biogeochemical plant-soil microbe feedback in response to climate warming. *Nature Climate Change* 3: 273-277.
- Breeuwer, A., Robroek, B.J.M., Limpens, J., Heijmans, M.M.P.D., Schouten, M.G.C. & Berendse, F. 2009. Decreased summer water table depth affects peatland vegetation. *Basic and Applied Ecology* 10: 330-339.
- Bokhorst, S., Huiskes, A., Aerts, R., Convey, P., Cooper, E.J., Dalen, L., Erschbamer, B., Gudmundsson, J., Hofgaard, A., Hollister, R.D., Johnstone, J. et al. 2013. Variable temperature effects of open top chamber at polar and alpine sites explained by irradiance and snow depth. *Global Change Biology* 19: 64-74.

- 457 Buttler, A., Grosvernier, Ph. & Matthey, Y. 1998. Development of *Sphagnum fallax* diaspores on  
458 bare peat with implications for the restoration of cut-over bogs. *Journal of Applied Ecology*  
459 35: 800-810.
- 460 Buttler, A. 1992. Permanent plot research in wet meadows and cutting experiment. *Vegetatio*  
461 103(2): 113-124.
- 462 Clymo, R.S. 1970. The Growth of Sphagnum: Methods of Measurement. *Journal of Ecology* 58(1):  
463 13-49.
- 464 Cullings, K.W. 1996. Single phylogenetic origin of ericoid mycorrhizae within the Ericaceae.  
465 *Canadian Journal of Botany* 74: 1896-1909.
- 466 Dabros, A., Fyles, J.W. & Strachan, I.B. 2010. Effects of open-top chambers on physical properties  
467 of air and soil at post-disturbance sites in northwestern Quebec. *Plant Soil* 333 : 203-218.
- 468 Delarue, F., Laggoun-Défarge, F., Buttler, A., Gogo, S., Jassey, V.E.J & Disnar, J.R. 2011. Effect  
469 of short-term ecosystem experimental warming on water extractable organic matter in an  
470 ombrotrophic *Sphagnum* peatland (Le Forbonnet, France). *Organic Geochemistry* 42: 1016-  
471 1024.
- 472 Delarue, F., Gogo, S., Buttler, A., Bragazza, L., Jassey, V., Bernard, G. & Laggoun-Défarge, F.  
473 2014. Indirect effects of experimental warming on dissolved organic carbon content in  
474 subsurface peat. *J. Soils Sediments* 14(11) : 1800-1805.
- 475 Delarue, F., Buttler, A., Bragazza, L., Grasset, L., Jassey, V.E.J., Gogo, S. & Laggoun, F. 2015.  
476 Experimental warming differentially affects microbial structure and activity in two  
477 contrasted moisture sites in a *Sphagnum*-dominated peatland. *Geoderma* (in press).
- 478 Dorrepaal, E., Aerts, R., Cornelissen, J.H.C. & Van Logtestijn, R.S.P. 2003. Summer warming and  
479 increased winter snow cover affect *Sphagnum fuscum* growth, structure and production in a  
480 sub-arctic bog. *Global Change Biology* 10: 93-104.
- 481 Dorrepaal, E., Cornelissen, J. H. C. & Aerts, R. 2007. Changing leaf litter feedbacks on plant  
482 production across contrasting sub-arctic peatland species and growth forms. *Oecologia* 151:  
483 251-261.
- 484 Dorrepaal, E., Aerts, R., Cornelissen, J.H.C., Van Logtestijn, R.S.P. & Callaghna, T.V. 2006.  
485 *Sphagnum* modifies climate-change impacts on subarctic vascular bog plants. *Functional*  
486 *Ecology* 20: 31-41.
- 487 Dorrepaal, E., Toet, S., van Logtestijn, R.S.P., Swart, E., van de Weg, M.J. Callaghan, T.V. &  
488 Aerts, R. 2009. Carbon respiration from subsurface peat accelerated by climate warming in  
489 the subarctic. *Nature* 460: 616-619.

- 490 Gogo S., Laggoun-Défarge F., Delarue F. & Lottier N., 2011. Invasion of a Sphagnum-peatland by  
491 Betula spp and Molinia caerulea impacts organic matter biochemistry. Implications for  
492 carbon and nutrient cycling. *Biogeochemistry* 106: 1, 53-69
- 493 Grosvernier, Ph., Matthey, Y. & Buttler, A. 1994 *Microclimate and physical properties of peat:*  
494 *new clues to the understanding of bog regeneration processes*. In: Restoration of Temperate  
495 Wetlands (Edited by B. D. Wheeler, S. C. Shaw, W. S. Fojt and R. A. Robertson), John  
496 Wiley & Sons, 435-450.
- 497 Gerdol, R., 1995. The growth dynamics of sphagnum based on field-measurements in a temperate  
498 bog and on laboratory cultures. *Journal of Ecology* 83: 431-437.
- 499
- 500 Hajek, T., Balance, S., Limpens, J., Zijlstra, M. & Verhoeven, J. T. A. 2011. Cell-wall  
501 polysaccharides play an important role in decay resistance of *Sphagnum* and actively  
502 depressed decomposition in vitro. *Biogeochemistry* 103: 45-57.
- 503 Heijmans, M.M.P.D., Klees, H. & Berendse, F. 2002. Competition between Sphagnum  
504 magellanicum and Eriophorum angustifolium as affected by raised CO<sub>2</sub> and increased N  
505 deposition. *Oikos* 97: 415–425.
- 506 Heijmans, M.M.P.D., Mauquoy, D, van Geel, B. and Berendse, F. 2008. Long-term effects of  
507 climate change on vegetation and carbon dynamics in peat bogs. *Journal of Vegetation*  
508 *Science* 19: 307-320.
- 509 Holden, J. 2005. Peatland hydrology and carbon release: why small-scale process matters. *Phil.*  
510 *Trans. R. Soc. A* 363: 2891–2913.
- 511 Hollister, R.D. & Webber, P.J. 2000. Biotic validation of small open-top chambers in a tundra  
512 ecosystem. *Global Change Biology* 6: 835-842.
- 513 Jacquemart, A.L. 1997. Vaccinium oxycoccus L. (Oxycoccus palustris Pers.) and Vaccinium  
514 microcarpum (Turez. ex Rupr.) Schmalh. (Oxycoccus microcarpum Turez. ex Rupr.).  
515 *Journal of Ecology* 85: 381–396.
- 516 Jacquemart, A.L. 1998. Biological Flora of the British Isles - Andromeda polifolia L. *Journal of*  
517 *Ecology* 86: 527-541.
- 518 Jassey, V.E.J., Chiapusio, G., Gilbert, D., Buttler, A. Toussaint, M-L., Binet, P. 2011. Experimental  
519 climate effect on seasonal variability of polyphenols/phenoloxidase interplay along a narrow  
520 fen-bog ecological gradient in *Sphagnum fallax*. *Global Change Biology* 17(9): 2945-2957.
- 521 Jassey V.E., Chiapusio G., Binet P., Buttler, A., Laggoun-Defarge, F., Delarue, F., Bernard, N.,  
522 Mitchell, E.A.D., Toussaint, M.L., Francez, A.J. & Gilbert, D. 2013. Above- and

- belowground linkages in Sphagnum peatland: climate warming affects plant-microbial interactions. *Global Change Biology*, 19, 811–823.
- Keuper, F., Dorrepaal, E., Van Bodegom, P.M., Aerts, R., Van Logtestijn, R.S.P., Callaghan, T.V. & Cornelissen, J.H.C. 2011. A Race for Space? How Sphagnum fuscum stabilizes vegetation composition during long-term climate manipulations. *Global Change Biology* 17(6): 2162-2171.
- Kuiper, J. J., Mooij, W. M., Bragazza, L. & Robroek, B.J.M. 2014. Plant functional types define magnitude of drought response in peatland CO<sub>2</sub> exchange. *Ecology* 95: 123–131.
- Laggoun-Défarge, F., Gilbert, D., Buttler, A., Epron, D., Francez, A.-J., Grasset, L., Guimbaud, C., Mitchell, E.A.D & Roy, J.-C. 2008. Effects of experimental warming on carbon sink function of a temperate pristine mire : the *PEATWARM* project. *Proceedings of the 13th International Peat Congress*, 599-602.
- Malmer, N., Svensson, B. & Wallen, B. 1994. Interactions between Sphagnum mosses and field layer vascular plants in the development of peat-forming systems. *Folia Geobotanica et Phytotaxonomica Praha* 29: 483-496
- Marion, G.M., Henry, G.H.R., Freckman, D.W., Johnstone, J., Jones, G., Jones, M.H., Levesque, E., Molau, U., Mølgaard, P., Parsons, A.N., Svoboda, J. & Virginia, R.A. 1997. Open-top designs for manipulating field temperature in high-latitude ecosystems. *Global Change Biology* 3: 20–32.
- Nijp, J.J., Limpens, J., Metselaar, K., van der Zee, S.E., Berendse, F. & Robroek, B.J. 2014. Can frequent precipitation moderate the impact of drought on peatmoss carbon uptake in northern peatlands? *New Phytologist* 203:70-80.
- Pinheiro, J.C. & Bates, D.M. 2000. *Linear mixed-effects models: basic concepts and examples*. Springer ISBN 978-1-4419-0318-1.
- R Core Team 2014. R: *A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- Robroek, B.J.M., Limpens, J., Breeuwer, A. & Schouten, M.G.C. 2007. Effect of water level and temperature on performance of four *Sphagnum* species. *Plant Ecology* 190: 97-107.
- Robroek, B.J.M., Heijboer, A., Jassey, V.E.J., Hefting, M.M., Rouwenhorst, T.G., Buttler, A. & Bragazza, L. 2013. Snow cover manipulation effects on microbial community structure and soil chemistry in a mountain bog. *Plant Soil* 369 (1-2) : 151-164.



- 555 Shaver, G.R. and Billings, W.D. 1977. Effect of day-length and temperature on root elongation in  
556 tundra graminoids. *Oecologia* 28: 57-65.
- 557 Soudani, K., Hmimina, G., Delpierre, N., Pontailier J.-Y., Aubinet, M., Bonal, D., Caquet, B., de  
558 Grandcourt, A., Burban. B., Flechard. C., Guyon D. et al. 2012. Ground-based Network of  
559 NDVI measurements for tracking temporal dynamics of canopy structure and vegetation  
560 phenology in different biomes. *Remote Sensing of Environment* 123: 234–245.
- 561 Sullivan, P.F. & Welker, J.M. 2005. Warming chambers stimulate early season growth of an arctic  
562 sedge: results of a minirhizotron field study. *Oecologia* 142: 616-626.
- 563 Titus, J.E., Wagner, D.J. & Stephens, M.D. 1983. Contrasting water relations of photosynthesis for  
564 two *Sphagnum* mosses. *Ecology* 64 (5): 1109-1115.
- 565 Turetsky, M.R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Frolking, S., McGuire, A.D. &  
566 Tuittila, E.S. 2012. The resilience and functional role of moss in boreal and arctic  
567 ecosystems. *New Phytologist* 196: 49-67.
- 568 Wallen, B. 1986. Above and below ground dry mass of the three main vascular plants on  
569 hummocks on a subarctic peat bog. *Oikos* 46: 51-56.
- 570 Walker, T.N., Ward, S.E., Ostle, N.J. & Bardgett, R.D. 2015. Contrasting growth responses of  
571 dominant peatland plants to warming and vegetation composition. *Oecologia*, DOI  
572 10.1007/s00442-015-3254-1.
- 573 Weltzin, J. F., Pastor, J., Harth, C., Bridgham, S. D., Updegraff, K., & Chapin, C. T. 2000.  
574 Response of bog and fen plant communities to warming and water-table manipulations.  
575 *Ecology* 81: 3464–3478.
- 576 Weltzin, J. F., Harth, C., Bridgham, S. D., Pastor, J. & Vonderharr, M. 2001. Production and  
577 microtopography of bog bryophytes: response to warming and water-table manipulations.  
578 *Oecologia* 128: 557–565.
- 579 Weltzin, J.F., Brigham, S.D., Pastor, J., Chen, J. & Harth, C. 2003. Potential effects of warming and  
580 drying on peatland plant community composition. *Global Change Biology*, 9, 141-151.
- 581 Yu, Z.C. 2011. Holocene carbon flux histories of the world's peatlands: global carbon-cycle  
582 implications. *Holocene* 21: 761-774.
- 583 Yan, W et al. 2014. Differentiating moss from higher plants is critical in studying the carbon cycle  
584 of the boreal biome. *Nature Communications* 5: 4270. DOI: 10.1038/ncomms5270  
585 |www.nature.com/naturecommunications.  
586  
587



588  
589

For Review Only

## Captions to figures

*Figure 1:* Changes of the relative frequencies of dominant plant species from 2008 to 2012 (no records in 2011) in the Forbonnet peatland (French Jura). Mean values (n=3) and standard error bars are given for dry (solid line) and wet (broken line) microhabitats, and for control (black line) and OTC (grey lines) plots. Significance of Linear Mixed Model is given for treatment effect (T) and for year effect (Y). Species that are not illustrated: *Carex pauciflora* (appeared in 2009 and then was only seen in one plot the years after), *Carex limosa* (appeared only in 3 plots in 2009), *Calluna vulgaris* (only in 2 different plots in different years), *Pinus rotundata* (only in one plot in one year).

*Figure 2:* Changes of the relative frequencies of species from April to September 2012 in the Forbonnet peatland (French Jura). Mean values (n=3) and standard error bars are given for dry (solid line) and wet (broken line) microhabitats, and for control (black line) and OTC (grey lines) plots. Significance of Linear Mixed Model is given for treatment (T) and month effect (M). Species that are not illustrated; with no significant differences: *Scheuchzeria palustris*, *Eriophorum vaginatum*, *Carex pauciflora*, *Carex rostrata*, *Drosera rotundifolia*, *Polytrichum strictum*; present only in one plot: *Calluna vulgaris*, *Carex pauciflora* and *Polytrichum strictum*.

*Figure 3:* Length increment (mm) of *Sphagnum* mosses in the different treatments (OTC vs control) and microhabitats (dry vs wet) in 2011 (a: May 13 – November 27, n=60) and 2012 (b: March 30 – November 6, n=120) in the Forbonnet peatland (French Jura). Vertical bars represent standard errors. Scales of both graphs are different. Significance levels of Linear Mixed Model are given for microhabitat (B) and treatment effect (T), and for their interaction (B x T).

*Figure 4:* Normalised Difference Vegetation Index (NDVI) at plot scale in the dry (a) and wet (b) microhabitats in year 2012 in the Forbonnet peatland (French Jura). Curves represent mean values (n=3) for control (black) and OTC (grey) plots. Significance codes for ANOVA's are (\*\*\*)  $p < 0.001$  ; (\*\*)  $p < 0.01$  ; (\*)  $p < 0.05$  ; (.)  $p < 0.1$ .

*Figure 1*

621  
622  
623  
624  
625  
626  
627

*Figure 2*

For Review Only

628

629 *Figure 3*

630

631

For Review Only

632  
633  
634  
635  
636  
637

*Figure 4*

For Review Only

Electronic appendices

Supporting Information to the paper Buttler et al. 2015. Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland. *Journal of Vegetation Science*. Appendix S1. Relationship between point-intercept frequency measures of vegetation and picture frequency analysis from the Forbonnet peatland (French Jura) for a: total vascular plants (Pearson  $r$ : 0.733,  $p < 0.01$ ), b: litter (Pearson  $r = 0.873$ ,  $p < 0.001$ ) and c: total *Sphagnum* and *Polytrichum strictum* (Pearson  $r = 0.942$ ,  $p < 0.001$ ).

The correlation between frequencies obtained by means of field point-intercepts and photographic analyses gives a good linear relation with vascular plants but shows an underestimation of mosses with the picture analysis. This is because leaves of vascular plants might hide the understory of mosses which than cannot be recorded in the one-layer photographic method, whereas in the field, the needle can hit more than one species or individual. This underestimation is mostly noticeable at intermediate moss cover, when vascular plants are most abundant. Conversely, the picture analysis overestimates litter when it is dense.











